arXiv:astro-ph/0212325v1 13 Dec 2002

SULPHUR CHEMISTRY IN THE ENVELOPES OF MASSIVE YOUNG STARS

F.F.S. van der Tak[1], **A.M.S. Boonman**[2], **R. Braakman**[2], **and E.F. van Dishoeck**[2]

[1]Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany
[2]Sterrewacht, Postbus 9513, 2300 RA Leiden, The Netherlands

Abstract

We present submillimeter observations of SO, SO_2, H_2S, H_2CS, OCS, NS and HCS^+ toward nine massive young stars. The outflow contributes ≈50% to the SO and SO_2 emission in $15-20''$ beams, more than for CS, where it is 10%. The SO_2 abundance increases from dark cloud levels in the outer envelope ($T < 100$ K) to levels seen in hot cores and shocks in the inner envelope ($T > 100$ K). Molecular abundances are consistent with a model of ice evaporation in an envelope with gradients in temperature and density for a chemical age of ~ 30000 yr. The high observed abundance of OCS, the fact that $T_{\rm ex}({\rm OCS}) \gg T_{\rm ex}({\rm H_2S})$, and the data on solid OCS and H_2S all suggest that the major sulphur carrier in grain mantles is OCS rather than H_2S. For most other sulphur-bearing molecules, the source-to-source abundance variations by factors of up to 10 do not correlate with previously established evolutionary trends in temperature tracers. These species probe the chemically inactive outer envelope. Our data set does not constrain the abundances of H_2S and SO in the inner envelope, which, together with SO_2, are required to use sulphur as a clock.

Key words: ISM: molecules – Molecular processes – Stars: Circumstellar matter; Stars: formation

1. Introduction

Spectral line surveys at submillimeter wavelengths have revealed considerable chemical differences between star-forming regions (see van Dishoeck 2001 for an overview). While these differences indicate activity, the dependence of molecular abundances on evolutionary state and physical parameters is poorly understood. Better insight into this relation would be valuable for probing the earliest, deeply embedded phases of star formation where diagnostics at optical and near-infrared wavelengths are unavailable. This is especially true for the formation of high-mass stars, for which the order in which phenomena occur is much less well understood than in the low-mass case, and for which the embedded phase is a significant fraction (≈10%) of the total lifetime of $\sim 10^6$ yr.



Figure 1. Examples of the spectra taken with the JCMT. The top and bottom frequency scales of the left panel are the two receiver sidebands; the right panel only shows lines from one sideband. Adopted from van der Tak et al. (2002).

Sulphur-bearing molecules are attractive as candidate tracers of early protostellar evolution (Hatchell et al 1998). Models of "hot core" chemistry (Charnley 1997), as well as of shock chemistry (Pineau des Forêts et al 1993) predict strong variations in the abundances of sulphur molecules on time scales of $\sim 10^4$ yr. To test these scenarios, this paper discusses submillimeter observations of sulphur-bearing molecules towards nine regions at the earliest stages of high-mass star formation, with luminosities $1 \times 10^4 - 2 \times 10^5$ $L_\odot$ at distances of $1-4$ kpc. This paper only summarizes our results; a complete description is given by van der Tak et al (2002).

Proceedings of the conference *"Chemistry as a Diagnostic of Star Formation,"* University of Waterloo, Canada, 21-23 August 2002 (C. L. Curry & M. Fich eds.).



Figure 2. Velocity profiles of the SO 6_6–5_5 line in the sources AFGL 2591 and W3 IRS5.



Figure 3. Distributions of the SO and SO_2 abundances in AFGL 2591 for chemical ages of 300, 3000, 30000 and 300000 yr (curves from bottom to top) after Doty et al. (2002). The $t \sim 3 \times 10^4$ yr model fits our data best.

2. Observations

Spectra in the 230 and 345 GHz bands were taken in 1995–1998 with the JCMT. The overview spectra in Fig. 1 show source-to-source differences in the relative and in the absolute strengths of lines of sulphur molecules. The line profiles (Fig. 2) reveal the presence of a narrow component, due to the envelope, and a broad one, due to the outflow. For SO and SO_2, the outflow contributes ≈50% of the line flux. This fraction is much larger than for CS, where it is only 10%. Lines of other molecules are too weak to separate the two components.

3. Results

To calculate molecular abundances, we have used the Monte Carlo program by Hogerheijde & van der Tak (2000), and models of the temperature and density structure of the sources (van der Tak et al 2000). Table 1 compares our results with abundances in other types of sources. The best overall match is with "hot cores".

For some sources, the detection of many SO_2 lines from a wide range of energy levels enables us to derive radial abundance profiles. Models with a constant SO_2 abundance of 10^{-9} fit the lines arising $\lesssim 100$ K above ground. However, the higher-excitation lines are underproduced, which indicates a 'jump' in the SO_2 abundance by a factor of ~100 at a temperature of ≈ 100 K. Infrared observations of SO_2 absorption by Keane et al (2001) also indicate abundances of $\sim 10^{-7}$ for the inner envelopes.

Table 1. Molecular abundances.

Source	CS 10^{-9}	H_2CS 10^{-10}	H_2S 10^{-9}	HCS^+ 10^{-10}	NS 10^{-11}	OCS 10^{-9}	SO 10^{-9}	SO_2 10^{-9}
W3 IRS5	5	3	9	0.2	...	0.5	10	10
W33A	5	13	8	2	2	20	1	1
AFGL 2136	4	3	<2	1	<0.5	...	1	0.5
AFGL 2591	10	3	20	2	0.5	10	10	2
S140 IRS1	5	7	4	5	1	<2	2	1
NGC 7538 IRS1	10	10	8	2	1	2	5	1
NGC 7538 IRS9	10	10	8	5	1	2	1	0.5
MonR2 IRS3	5	7	<4	...	...	...	5	1
NGC 6334 IRS1	10	7	10	2	10	50	2	2
Hot cores[a]	8	6	9	...	80	5	4	20
PDRs[b]	20	...	6	...	...	...	9	0.1
Dark clouds[c]	1	6	0.8	0.6	...	2	20	4
Shocks[d]	4	8	4000	0.1	...	10	200	100

[a] Source average from Hatchell et al (1998) [b] Orion Bar: Jansen et al (1995)
[c] L 134N: Ohishi et al (1992) [d] Orion Plateau: Sutton et al (1995); Minh et al (1990)

4. CHEMICAL IMPLICATIONS

Most hot core models only consider single values for the temperature and density, while for our sources, these parameters vary strongly along the line of sight. Using the UMIST database, Doty et al (2002) calculated the time-dependent gas-phase chemistry for the temperature and density structure of AFGL 2591. The model has most sulphur initially in S at $T < 100$ K and in H_2S at $T > 100$ K, to mimic the effect of ice evaporation. The resulting 'jump' in the abundances at $r = 2 \times 10^{16}$ cm is consistent with our observations of SO_2. See Fig. 3 and Doty (this volume).

The model reproduces most of our observed abundances, and also the increase of the SO_2 abundance at $T \gtrsim 100$ K. However, OCS is underproduced by factors of 10–100, suggesting that most of the sulphur in the grain mantles is in the form of OCS. This idea is supported by the high excitation temperature of OCS (100 K), much higher than the 25 K found for H_2S. In addition, solid OCS has been detected in the infrared towards W33A, while H_2S has not.

The current data set is not suitable to trace chemical evolution within our sample. The CS, H_2CS, HCS^+ and NS molecules mainly probe the chemically inactive outer envelope ($T < 100$ K). While OCS, SO and SO_2 show larger source-to-source variations, only for SO_2, the data allow to determine the abundance in the inner envelope ($T < 100$ K) where ices are evaporating. The situation for H_2S is unclear: models predict strong abundance variations, but the present data do not trace the warm gas. Future observations of high-excitation lines of H_2S, SO and OCS could exploit the full potential of sulphur as chemical clock.

REFERENCES

Charnley, S. B. 1997, ApJ, 481, 396

Doty, S. D., van Dishoeck, E. F., van der Tak, F. F. S., & Boonman, A. M. S. 2002, A&A, 389, 446

Hatchell, J., Thompson, M. A., Millar, T. J., & MacDonald, G. H. 1998, A&A, 338, 713

Hogerheijde, M. & van der Tak, F. 2000, A&A, 362, 697

Jansen, D. J., Spaans, M., Hogerheijde, M. R., & van Dishoeck, E. F. 1995, A&A, 303, 541

Keane, J. V., Boonman, A. M. S., Tielens, A. G. G. M., & van Dishoeck, E. F. 2001, A&A, 376, L5

Minh, Y. C., Irvine, W. M., McGonagle, D., & Ziurys, L. M. 1990, ApJ, 360, 136

Ohishi, M., Irvine, W. M., & Kaifu, N. 1992, in IAU Symp. 150: Astrochemistry of Cosmic Phenomena, 171

Pineau des Forêts, G., Roueff, E., Schilke, P., & Flower, D. R. 1993, MNRAS, 262, 915

Sutton, E., et al 1995, ApJS, 97, 455

van der Tak, F. F. S., Boonman, A. M. S., Braakman, R., & van Dishoeck, E. F. 2002, A&A, submitted

van der Tak, F. F. S., van Dishoeck, E. F., Evans, N. J., & Blake, G. A. 2000, ApJ, 537, 283

van Dishoeck, E. F. 2001, The Promise of the HSO, eds. G.L. Pilbratt et al. ESA-SP, 460, 185